

Practical guide to analytic loop integration

Exercise Sheet 1

<https://yannickulrich.gitlab.io/loop-integration>

Issued: 13 May 2022

Exercise 1: Γ functions

Use our basic integral

$$\int_0^\infty dx x^\alpha (a + bx)^\beta = \frac{\Gamma(1 + \alpha)\Gamma(-1 - \alpha - \beta)}{\Gamma(-\beta)} \frac{a^{1+\alpha+\beta}}{b^{1+\alpha}} \quad (1)$$

to calculate





a) $I_1(\alpha, \beta, \gamma, \delta) = \int dx_1 dx_2 dx_3 \delta(\dots) x_1^\alpha x_2^\beta x_3^\gamma (x_2 x_3 + x_1 x_2 + x_1 x_3)^\delta$

b) $I_2(\alpha, \beta, \gamma) = \int dx_1 dx_2 dx_3 dx_4 \delta(\dots) x_1^\alpha (x_2 x_3 + x_2 x_4 + x_3 x_4)^\beta (x_1 x_3 + x_2 x_3 + x_1 x_4 + x_2 x_4 + x_3 x_4)^\gamma$

c) $I_2(-2\epsilon, -1 - 2\epsilon, -1 + 3\epsilon)$ up to $\mathcal{O}(\epsilon^0)$

Exercise 2: Graph polynomials

Find the graph polynomials and calculate the loop integrals



Dashed lines represent massless propagators and solid lines massive ones.

Exercise 3: Sunset integral

For general masses the sunset integral of 2b cannot be solved at two-loop. However, in the example above with massless internal lines and $p^2 \neq 0$ it was rather easy.

- a) This is also true at three loop. Show that

$$I_{\ell=3} = \text{---} \langle \text{Sunset Diagram} \rangle \text{---} = (-p^2)^{2-3\epsilon} \frac{\Gamma(1-\epsilon)^7 \Gamma(-2+3\epsilon)}{\Gamma(4-4\epsilon)}.$$

- b) Find a recurrence relation for the ℓ -loop graph polynomials \mathcal{F}_ℓ and \mathcal{U}_ℓ as a function of $\mathcal{F}_{\ell-1}$ and $\mathcal{U}_{\ell-1}$. Then construct the ℓ -loop sunset I_ℓ , integrate over one Feynman parameter, and relate the result to $I_{\ell-1}$. Finally, solve the resulting recurrence relation to obtain

$$I_\ell = \text{---} \langle \text{\ell-loop Sunset Diagram} \rangle \text{---} = (-1)^{\ell+1} (-p^2)^{\ell-1-\ell\epsilon} \frac{\Gamma(1-\epsilon)^{2\ell+1} \Gamma(1-\ell(1-\epsilon))}{\Gamma((1+\ell)(1-\epsilon))}$$

- c) Write code to calculate I_ℓ using only the algorithm for \mathcal{U} and \mathcal{F} and the master formula. (*for the adventurous*)
- d) Use the optical theorem to relate $\Im I_\ell$ to the phase space volume for a $1 \rightarrow n$ decay

$$\int d\text{PS}^{1 \rightarrow n} = \frac{s^{n-2}}{2(4\pi)^{2n-3} \Gamma(n) \Gamma(n-1)}.$$